

# Three Faces of the Delta Conjecture

J. Haglund  
University of Pennsylvania

March 30, 2019

# The Algebraic Side

Let  $X_n = \{x_1, \dots, x_n\}$ ,  $Y_n = \{y_1, \dots, y_n\}$  be sets of variables. Let

$$DR_n = \mathbb{C}[X_n, Y_n] / \left\{ \sum_i x_i^a y_i^b : a, b \geq 0, a + b > 0 \right\}$$

be the ring of diagonal coinvariants.  $S_n$  acts “diagonally” on  $DR_n$  by permuting the  $X$  and  $Y$  variables in the same way.

Example:  $n = 2$

Cosets  $\{1, x_1, y_1\}$  form a basis for  $DR_2$ , so  $\text{Hilb}(DR_2) = 1 + q + t$ .

The identity in  $S_2$  acts by fixing all the cosets, while  $\sigma = (12)$  fixes 1 and sends  $\{x_1, y_1\}$  to  $\{x_2, y_2\}$ . Since  $x_1 + x_2 = 0 = y_1 + y_2$ ,  $x_2 = -x_1, y_2 = -y_1$ . Hence the coset 1 corresponds to the trivial character, while  $x_1, y_1$  correspond to the sign character, and the bigraded character of  $DH_2$  is  $s_2 + (q + t)s_{1,1}$ .

# The Symmetric Function Side

Let  $\Delta'_f$  be a linear operator defined via

$$\Delta'_f \tilde{H}_\mu(X; q, t) = f[B_\mu - 1] \tilde{H}_\mu(X; q, t),$$

where  $B_\mu = \sum_{s \in \mu} q^{\text{coarm}(s)} t^{\text{coleg}(s)}$ . For example

$$B_{3,2} = 1 + q + q^2 + t + tq.$$

Haiman proved that the bigraded character of  $\text{DR}_n$  under the diagonal action is given by

$$\Delta'_{e_{n-1}} e_n(X) = \sum_{\mu \vdash n} \frac{T_\mu \tilde{H}_\mu(X; q, t) M B_\mu \prod'_{s \in \mu} (1 - q^{\text{coarm}(s)}) (1 - t^{\text{coleg}(s)})}{\prod_{s \in \mu} (t^{\text{leg}(s)} - q^{\text{arm}(s)+1}) (q^{\text{arm}(s)} - t^{\text{leg}(s)+1})}$$

where  $M = (1 - q)(1 - t)$  and  $T_\mu = t^{n(\mu)} q^{n(\mu')}$ , with  $n(\mu) = \sum_i (i - 1) \mu_i$ .

# The Combinatorial Side

Given a Dyck path  $\pi$  and a word parking function  $P$  (a filling of the squares just to the right of North steps of  $\pi$  with cars, i.e. integers between 1 and  $n$ , strictly increasing up columns), let  $a_i$  be the number of area squares in the  $i$ th row (from the bottom). Cars in rows  $(i, j)$  with  $i < j$  form an inversion pair if either  $a_i = a_j$  and  $car_i < car_j$ , or  $a_i = a_j + 1$  and  $car_i > car_j$ . Let  $d_i$  be the number of inversion pairs  $(i, j)$  with  $i < j$ . Furthermore, we call a car at the bottom of a column a *valley*, and say the valley is *moveable* if, when we slide the car one square to the left, the result is still a word parking function, i.e. we still have strict decrease down columns. For example, in Figure 1, cars 1, 2 and 8 (in rows 5, 6 and 8) are moveable, but cars 4 and 3 in rows 1 and 2 are not.




Figure: A word parking function with area = 6. There are  $\text{div}v(i, j)$ -row pairs  $(7, 8)$ ,  $(5, 7)$ ,  $(5, 8)$ ,  $(4, 5)$ ,  $(4, 7)$ ,  $(3, 6)$ ,  $(3, 8)$ , so  $\text{div}v = 7$ . The total weight is  $x_1 x_2 x_3 x_4^2 x_7 x_8^2 q^7 t^6$ .

## Theorem (Carlsson-Mellit, 2015)

$$\Delta'_{e_{n-1}} e_n = \sum_{P \in \text{WP}(n)} q^{\text{dinv}(P)} t^{\text{area}(P)} x^P.$$

where the sum is over all word parking functions  $P$  on  $n$  cars.

Still Open: Find a combinatorial expression for the Schur expansion of the right-hand-side above.

## Corollary (Conjectured by H., Loehr in 2002)

$$\text{Hilb}(\text{DR}_n) = \sum_{\sigma \in \mathcal{S}_n} t^{\text{maj}(\sigma)} \prod_{i=1}^{n-1} [w_i(\sigma)]_q.$$

Let  $w_i(\sigma)$  equal the number of  $w_j$  which are in  $\sigma_i$ 's run and larger than  $\sigma_i$ , or in the next run to the right and smaller than  $\sigma_i$ .

### Example

$$\sigma = 25713846 \rightarrow 257|138|46|0$$
$$(w_1, w_2, \dots, w_8) = (3, 3, 2, 2, 1, 2, 2, 1).$$

## Theorem (Carlsson-Oblomkov, 2018)

A monomial basis for  $DR_n$  is given by a certain family of cosets, one for each  $\sigma \in S_n$ . The contribution to  $\text{Hilb}(DR_n)$  of monomials associated to  $\sigma$  is  $t^{\text{maj}(\sigma)} \prod_{i=1}^{n-1} [w_i(\sigma)]_q$ .

## Examples

$$\sigma = 25713846 \rightarrow y_1 y_2 y_3 \times y_1 y_2 y_3 y_4 y_5 y_6$$

$$(1 + x_2 + x_2^2)(1 + x_5 + x_5^2)(1 + x_7)(1 + x_1)(1 + x_8)(1 + x_4)$$

Set all  $x_i = 0$ ;  $\sum_{\sigma \in S_n} \prod_{k \in \text{Des}} y_k \rightarrow$  Garsia-Stanton basis

Set all  $y_i = 0$ ;  $\sigma = (12 \cdots n) : (w_1, w_2, \dots) = (n, n-1, \dots) \rightarrow$   
 $(1 + x_1 + \dots + x_1^{n-1}) \cdots (1 + x_{n-2} + x_{n-2}^2)(1 + x_{n-1}) \rightarrow$  Artin basis.



					8		1
					2		0
			6				2
		5					2
		4					1
7							2
3							1
1							0

## The Delta Conjecture (H., Remmel, Wilson, 2015)

$$\begin{aligned}
 \Delta'_{e_{k-1}} e_n &= \sum_{P \in \text{WP}(n)} q^{\text{dinv}(P)} t^{\text{area}(P)} \prod_{a_i > a_{i-1}} (1 + z/t^{a_i}) \Big|_{z^{n-k}} \\
 &= \sum_{P \in \text{WP}(n)} q^{\text{dinv}(P)} t^{\text{area}(P)} \prod_{\text{movable valleys}} (1 + z/q^{d_i+1}) \Big|_{z^{n-k}}
 \end{aligned}$$

Let  $\Pi$  be an ordered set partition of  $\{1, 2, \dots, n\}$ , and let  $\sigma = \sigma(\Pi)$  be the ordering of the blocks of  $\Pi$  which minimizes  $\text{maj}$ . For example, if  $\Pi = \{\{2, 3, 5\}, \{1, 6, 7, 9\}, \{4, 8\}\}$ , then  $\sigma(\Pi) = 235679148$ , and  $\text{minimaj}(\Pi) = \text{maj}(\sigma) = 6$ . Next form  $\sigma^*$  by marking every number which is not leftmost (in  $\text{minimaj}$  order) from its block;

$$\sigma^* = 23^*5^*67^*9^*1^*48^*.$$

Now construct the vector  $(w_1(\Pi), w_2(\Pi), \dots)$  by first isolating the unmarked elements of  $\sigma^*$ , map them to a permutation, and apply previous rule:

$$264 \rightarrow 132 \rightarrow 13|2|0 \rightarrow (1, 1, 1).$$

For marked elements  $\sigma_i^*$ ,  $w_i$  equals the number of unmarked elements smaller than  $\sigma_i$  in its run plus the number of unmarked elements which are larger in the previous run.

$$\sigma^* = \{23^*5^*\}\{67^*9^*1^*\}\{48^*\} \rightarrow (1, 1, 1, 1, 2, 2, 2, 1, 1).$$

## Theorem H.-Sergel, 2018

$$\sum_{P \in \text{PF}(n)} q^{\text{dinv}(P)} t^{\text{area}(P)} \prod_{\text{movable valleys}} (1 + z/q^{d_i+1}) \Big|_{z^{n-k}} = \sum_{\substack{\Pi \\ k \text{ blocks}}} t^{\text{minimaj}(\Pi)} \prod_{i=1}^n [w_i(\Pi)]_q.$$

Open Question: Is there an analogue involving the rise version of the Delta Conjecture?

# A module for the Delta Conjecture

M. Zabrocki has recently introduced a module whose bigraded character is conjecturally equal to the combinatorial and symmetric function sides of the Delta Conjecture. Let  $\Theta_n = \{\theta_1, \dots, \theta_n\}$  be a set of anticommuting variables, i.e.  $\theta_i\theta_j = -\theta_j\theta_i$ ,  $1 \leq i < j \leq n$ . Note this implies  $\theta_i^2 = 0$ . Let  $X_n, Y_n$  be two sets of commuting variables, which also commute with the  $\theta_i$ . Set

$$\text{TR}_n = \mathbb{C}[X_n, Y_n, \Theta_n] / \left\{ \sum_i x_i^a y_i^b \theta_i^c : a, b, c \geq 0, a + b + c > 0, c \leq 1 \right\}.$$

$S_n$  acts on  $\text{TR}_n$  diagonally by permuting the  $x_i, y_i, \theta_i$  in the same way. Then Zabrocki conjectures that the tri-graded character of this action is given by

$$\sum_{k=1}^n z^{n-k} \Delta'_{e_{k-1}} e_n,$$

where  $q, t$  give the grading in the  $x$  and  $y$  variables and  $z$  the grading in the  $\theta$  variables.